

Entropic uncertainty relations for incomplete sets of mutually unbiased observables

Adam Azarchs, California Institute of Technology

February 1, 2008

Abstract

Entropic uncertainty relations, based on sums of entropies of probability distributions arising from different measurements on a given pure state, can be seen as a generalization of the Heisenberg uncertainty relation that is in many cases a more useful way to quantify incompatibility between observables. Of particular interest are relationships between ‘mutually unbiased’ observables, which are maximally incompatible. Lower bounds on the sum of entropies for sets of two such observables, and for complete sets of observables within a space of given dimension, have been found. This paper explores relations in the intermediate regime of large, but far from complete, sets of unbiased observables.

1 Background and prior work

Entropic uncertainty relations are analogous to the Heisenberg uncertainty principle, but phrased in terms of the entropy of incompatible measurements rather than the variances. The entropic form is preferable to the original Heisenberg formulation in many situations. The entropy of measuring a state $|\psi\rangle$ with respect to basis $\{|a_i^{(k)}\rangle\}_{i=1}^N$ is defined for these purposes as

$$H_k = - \sum_{i=1}^N p_i^{(k)} \log p_i^{(k)},$$

where N is the dimension and $p_i^{(k)}$ is the probability of the measurement yielding outcome i . That is,

$$p_i^{(k)} = |\langle a_i^{(k)} | \psi \rangle|^2.$$

David Deutsch [1] found a relationship

$$H_a + H_b \geq -2 \log \frac{1}{2}(1 + c),$$

where

$$c = \max_{j,k} |\langle a_j | b_k \rangle|.$$

Maassen and Uffink [2] improved this bound to

$$H_a + H_b \geq -2 \log c. \quad (1)$$

This later relationship is much stronger since $c \ll 1$ for large N .

Of particular interest are so-called mutually unbiased observables (or bases), defined as those where

$$|\langle a_i^{(k)} | a_j^{(l)} \rangle|^2 = 1/N$$

for all $k \neq l$. It has been shown [3] that at least for prime N there exist sets of $N + 1$ such observables. Sánchez [4] found that for such sets, one could write

$$\sum_{k=1}^{N+1} H_k \geq (N + 1) \log \frac{1}{2} (N + 1). \quad (2)$$

This is substantially better than the bound derived by breaking up the set pair-wise and using equation 1,

$$\sum_{k=1}^{N+1} H_k \geq \frac{1}{2} (N + 1) \log N. \quad (3)$$

For several interesting applications, such as encryption, a relationship like (2) summing only over the incomplete set of M observables, would be useful. Weak lower bounds can be formed by modification of (3) to

$$\sum_{k=1}^M H_k \geq \frac{1}{2} M \log N. \quad (4)$$

However, this quickly becomes a poor bound as M increases. Similarly, for M approaching $N + 1$ we can take (2) and subtract the upper bound on the entropy of the observables that are being left out, $(N + 1 - M) \log N$, which yields

$$\sum_{k=1}^M H_k \geq (N + 1) \log \frac{N + 1}{2N} + M \log N. \quad (5)$$

However for $M \ll N$ this is actually negative, so useless in that regime. The regime where N is very large, and M is also large but $\ll N$, is quite interesting to ‘locking’ correlations [5]. A relation useful in the domain around $M \sim N / \log N$ is derived in this paper.

2 Result for intermediate domain

Larsen [6] has derived the equality relationship

$$\sum_{k=1}^{N+1} \pi_k = \Pi + 1$$




Figure 1: Equation (9) vs. M for $N = 1009$. The dashed lines represent the weaker lower bounds (4) and (5).

where the π_k s are the ‘purities,’

$$\pi_k = \sum_{i=1}^N [p_i^{(k)}]^2,$$

and Π is the purity $\text{Tr}(\rho^2)$ of the state ρ . Clearly, $\frac{1}{N} \leq \pi_k \leq 1$, and similarly for Π . Then subtracting the excess we can see that

$$\sum_{k=1}^M \pi_k \leq \Pi + 1 - \frac{N+1-M}{N} \leq \frac{N-1+M}{N}. \quad (6)$$

Since the product of a set of positive numbers whose sum is constrained is maximized when the numbers are all equal, we can use (6) to write

$$\prod_{k=1}^M \pi_k \leq \left(\frac{N-1+M}{NM} \right)^M. \quad (7)$$

Because $H_k \geq -\ln \pi_k$, [2] this gives us the relation

$$\sum_{k=1}^M H_k \geq M \log \frac{NM}{N-1+M}. \quad (8)$$

Sánchez [7] showed that his bound (2) for complete sets, where the dimension N is even, cannot be tight as the bound $H_k \geq -\log \pi_k$ can only be tight for $\pi_k = 1/m$, $m \in \mathbb{N}$. For $\frac{1}{m} \leq \pi_k \leq \frac{1}{m-1}$, $z \in \mathbb{N}$ he derived [8] the convex inequality

$$H_k \geq \log m - (m-1)(m\pi_k - 1) \log \frac{m}{m-1}$$

This can be applied to strengthen relationship (8), yielding

$$\sum_{k=1}^M H_k \geq M \left[\log m - (m-1) \left(m \frac{N+M-1}{NM} - 1 \right) \log \frac{m}{m-1} \right] \quad (9)$$

where

$$m = \text{ceil} \left(\frac{NM}{N+M-1} \right).$$

Figure 1 plots this bound with respect to M .

Relations (8) and (9), for $M \sim N/\log N$ (near the intersection of (4) and (5)) provide a significantly improved bounds. The behavior at the end points $M = 1$ and $M = N+1$ is correct as well. Relation (8) is only marginally weaker, and then only for $NM/(N+M-1) \notin \mathbb{N}$, and has a somewhat more convenient form.

References

- [1] David Deutsch. Uncertainty in Quantum Measurements. *Physical Review Letters*, 50(9):631–633, February 1982.
- [2] Hans Maassen and J. B. M. Uffink. Generalized entropic uncertainty relations. *Physical Review Letters*, 60(12):1103–1106, March 1988.
- [3] I D Ivonovic. Geometrical description of quantal state determination. *Journal of Physics A: Mathematical and General*, 14(12):3241–3245, December 1981.
- [4] Jorge Sánchez. Entropic uncertainty and certainty relations for complementary observables. *Physics Letters A*, 173:233–239, 1993.
- [5] David P. DiVincenzo, Michal Horodecki, Debbie W. Leung, John A. Smolin, and Barbara M. Terhal. Locking classical correlation in quantum states. *Physical Review Letters*, 92, 2004.
- [6] Ulf Larsen. Superspace geometry: the exact uncertainty relationship between complementary aspects. *Journal of Physics A: Mathematical and General*, 23:1041–1061, 1990.
- [7] Jorge Sánchez. States of minimal joint uncertainty for complementary observables in three-dimensional hilbert space. *Journal of Physics A: Mathematical and General*, 27:843–846, 1994.
- [8] Jorge Sánchez. Improved bounds in the entropic uncertainty and certainty relations for complementary observables. *Physics Letters A*, 201:125–131, May 1995.

Acknowledgements

I would like to thank my mentors Drs. John Preskill and Patrick Hayden for their direct support in this project. I would also thank Rohit Thomas and Peter Knophf for their help with untangling some nasty math, and the donor for the Victor Neher named SURF fellowship for financial support.